

On the variety of Borels in relative position $\vec{w}$

David Joyner and Pablo Lejarraga

1-31-2003

Abstract

Let G be a connected semi-simple group defined over and algebraically closed field, T a fixed Cartan, B a fixed Borel containing T , S a set of simple reflections associated to the simple positive roots corresponding to (T, B) , and let $\mathcal{B} \cong G/B$ denote the Borel variety. For any $s_i \in S$, $1 \leq i \leq n$, let $\overline{O}(s_1, \dots, s_n) = \{(B_0, \dots, B_n) \in \mathcal{B}^{n+1} \mid (B_{i-1}, B_i) \in \overline{O}(s_i), 1 \leq i \leq n\}$, where $O(s)$ denotes the subvariety of pairs of Borels in $\mathcal{B}^2$ in relative position s . We show that such varieties are smooth and indicate why this result is, in one sense, best possible. Our main results assume that k has characteristic 0.

Contents

1	Main results	3
2	Definitions and lemmas	5
3	The smoothness result	10
4	Remarks on $\mathbb{P}^1$-bundles	11
5	Deligne-Lusztig varieties	12
6	Appendix: Technical background	13
6.1	Standard properties of fibers	13
6.2	Main lemma	14

Let G be a connected semi-simple group defined over and algebraically closed field, T a fixed Cartan, B a fixed Borel containing T , S a set of simple reflections associated to the simple positive roots corresponding to (T, B) , W the Weyl group of T , and let $\mathcal{B} \cong G/B$ denote the Borel variety. For $w_i \in W$, let $\vec{w} = (w_1, \dots, w_n)$ and $\overline{O}(\vec{w}) = \{(B_0, \dots, B_n) \in \mathcal{B}^{n+1} \mid (B_{i-1}, B_i) \in \overline{O(w_i)}, 1 \leq i \leq n\}$, where $O(w)$ denotes the subvariety of pairs of Borels in $\mathcal{B}^2$ in relative position w (defined below) and $\overline{O(w)}$ denotes Zariski closure of $O(w)$ in $\mathcal{B} \times \mathcal{B}$. We call such a variety the **closed variety of Borels of relative position $\vec{w}$** .

Such varieties have appeared in many works. We mention, for example, [Han1], [DL], and [Spr1]. In special cases, they are known to be non-singular. Let $\vec{s} = (s_1, \dots, s_n) \in S^n$. We prove that these varieties $\overline{O}(\vec{s})$ are non-singular. Although this seems to be a simple result, we did not succeed in finding a simple proof. The problem lies in the distinction between reduced fibers and schematic fibers in the statement of a theorem of Mumford (Theorem 3' in §III.10 of [Mum]) which we use at one point in the proof. Most of the technicalities, which ultimately rely on results of Grothendieck [EGA], have been pushed into §6.

We also prove an analogous smoothness result for the corresponding Deligne-Lusztig varieties, under a technical assumption. In some cases, such varieties are known to have a large number of rational points, so there is the hope that such results will be useful for constructing error-correcting codes having “good” parameters [Han2].

Finally, we indicate why our smoothness result is best possible in some sense. Consider the projection $\pi_w : \overline{O}(w) \rightarrow \overline{O}(\emptyset) = \mathcal{B}$, where $w \in W$ is *not* a simple reflection. In general, it is known that $\overline{O}(w)$ is singular (this is the *raison d’être* for the Demazure-Hansen resolution [DL], §9.1) and of course $\overline{O}(\emptyset)$ (the Borel variety) is smooth. Therefore, by Lemma 6.1 below, π_w cannot be smooth. As a consequence of this type of reasoning, our smoothness result cannot be generalized to the analogous projection $\overline{O}(w_1, \dots, w_n) \rightarrow \overline{O}(w_1, \dots, w_{n-1})$, $w_i \in W$.

Acknowledgement: We thank S. Hansen for useful correspondence on [Han1], [Han2].

1 Main results

Let G be a connected semisimple group defined over an algebraically closed field k with maximal torus T contained in a Borel B . This is given the structure of a Coxeter group generated by the reflections s_α associated to the simple positive roots $\alpha \in \Delta$ of (T, B) . We identify G with the group of k -valued points of G .

We shall denote conjugation of group elements by ${}^gx = gxg^{-1}$. Representing $w \in W$ by $\dot{w} \in N_G(T)$, define

$${}^wB' = \dot{w}B'\dot{w}^{-1},$$

for any Borel B' of G . The Borel subgroups belonging to the set $\{{}^wB \mid w \in W\}$ are precisely the Borel subgroups of G which contain T .

Let $\mathcal{B} \cong G/B$ denote the Borel variety.

We say that $B', B'' \in \mathcal{B}$ are in **position** w if (B', B'') belongs to the G -orbit of $(B, {}^wB)$, for $w \in W$. We denote this orbit by

$$O(w) = \{(B', B'') \in \mathcal{B}^2 \mid B', B'' \text{ are in position } w\}.$$

In particular, $O(1) \cong \mathcal{B}$ is the diagonal in $\mathcal{B}^2$. This orbit $O(w)$ is open in its Zariski closure $\overline{O(w)} \subset \mathcal{B} \times \mathcal{B}$ (this follows from a much more general result in [St], using the fact that $\mathcal{B}$ has a transitive G -action). It is known that

$$\overline{O(w)} = \cup_{w' \leq w} O(w'),$$

where $\leq$ denotes the Bruhat ordering associated to (T, B) ([Spr1], §10.2.13).

Let $\vec{w} = (w_1, \dots, w_n) \in W^n$ and let

$$O(\vec{w}) = \{(B_0, \dots, B_n) \in \mathcal{B}^{n+1} \mid (B_{i-1}, B_i) \in O(w_i), 1 \leq i \leq n\}.$$

We call this the **variety of Borels of relative position** $\vec{w}$.

We say an algebraic group G **operates** on a variety X if there is a morphism $\phi : G \times X \rightarrow X$ satisfying $\phi(1, x) = x$ for all $x \in X$ (which may or may be compatible with the group operation). We say G **acts** on X if it operates on X in a way compatible with the group operation. For notational simplicity, we usually suppress the ϕ from the notation and use juxtaposition to denote an operation.

A basic fact about this variety is the following.

Lemma 1.1. *Let $\vec{s} = (s_1, \dots, s_n) \in S^n$. The variety $O(\vec{s})$ is the orbit of a connected algebraic group operation.*

This is proven in §2.

Let

$$\overline{O}(w_1, \dots, w_n) = \{(B_0, \dots, B_n) \in \mathcal{B}^{n+1} \mid (B_{i-1}, B_i) \in \overline{O}(w_i), 1 \leq i \leq n\}.$$

This variety is closed in $\mathcal{B}^{n+1}$. In Lemma 3.1, we will see that this may be written as an iterated fiber product of the $\overline{O}(w_i)$'s. The projection morphism

$$\pi_n : \overline{O}(w_1, \dots, w_n) \rightarrow \overline{O}(w_1, \dots, w_{n-1}),$$

is defined¹ by ignoring the last “coordinate”. We show in §3 that π_n is flat. However, in the case when the w_i are simple reflections, much more is true.

Proposition 1.2. *Assume that k has characteristic 0. Let $\vec{s} = (s_1, \dots, s_n) \in S^n$. For $n \geq 1$, $\overline{O}(s_1, \dots, s_n)$ is non-singular, and π_n is smooth.*

This is proven in §3. We show in §2 and §6 that each schematic fiber of π_n is $\cong \mathbb{P}^1$ (for $\text{char}(k) = 0$ - in fact, this result is needed to prove the above proposition and is the reason for the characteristic 0 hypothesis there). We believe that these results hold even in characteristic $p > 0$.

Corollary 1.3. *If $w = s_1 \dots s_n$ is a minimal expression then the projection map $(B_0, \dots, B_n) \mapsto (B_0, B_n)$ induces a surjection the morphism $\pi : \overline{O}(s_1, \dots, s_n) \rightarrow \overline{O}(w)$ such that*

$$\pi : \pi^{-1}(O(w)) \rightarrow O(w) \tag{1}$$

is an isomorphism. In other words, π is a resolution of singularities of $\overline{O}(w)$.

This is well-known (see, for example, §9 in Deligne-Lusztig [DL] or §10.2 in Springer [Spr1]). If, in addition, the $s_1, \dots, s_n$ are distinct then this resolution is an isomorphism [Han1].

Question 1.4. *Is*

$$\overline{O}(s_1, \dots, s_n) \xrightarrow{\pi_n} \overline{O}(s_1, \dots, s_{n-1})$$

a $\mathbb{P}^1$ -bundle?

¹If $n = 1$, we interpret $\overline{O}(w_1, \dots, w_0) = \overline{O}(\emptyset)$ as $\mathcal{B}$.

Some remarks on this are given in §4.

Let $\vec{s} = (s_1, \dots, s_n) \in S^n$ be a list of simple reflections, with respect to a fixed Borel B in a connected semisimple group (functor) G defined over a finite field $\mathbb{F}_q$. Let $F : G \rightarrow G$ denote the Frobenius morphism. Following Hansen [Han1], let

$$\overline{X}(\vec{s}) = \{({}^{g_0}B, \dots, {}^{g_n}B) \in \mathcal{B}^{n+1} \mid g_k^{-1}g_{k+1} \in B \cup Bs_{k+1}B, \text{ for } 0 \leq k < n, g_n \in F(g_0)\}.$$

Here we shall call $\overline{X}(\vec{s})$ a **Deligne-Lusztig variety**.

Corollary 1.5. *Assume that Proposition 1.2 holds for $k = \overline{\mathbb{F}_q}$. $\overline{X}(\vec{s})$ is non-singular.*

This is proven in §5.

2 Definitions and lemmas

We begin with some simple observations on terminology. We call $\overline{O}(s_1, \dots, s_n)$ the closed variety of Borels of relative position $\vec{w}$. On the other hand, the closure of the variety of Borels of relative position $\vec{w}$ is $\overline{O(s_1, \dots, s_n)}$. How are these related?

Lemma 2.1. $\overline{O}(s_1, \dots, s_n) = \overline{O(s_1, \dots, s_n)}$.

Remark 2.2. *It is clear that $\overline{O}(s_1, \dots, s_n)$ is closed in $\mathcal{B}^{n+1}$ since it is defined by closed conditions. This implies $\overline{O(s_1, \dots, s_n)} \subset \overline{O}(s_1, \dots, s_n)$. By Lemma 1.1 above and the proof of Lemma 2.3.3(i) in [Spr2], $O(s_1, \dots, s_n)$ contains a set which is open in $\overline{O(s_1, \dots, s_n)}$. Observe that $O(s_1, \dots, s_n)$ is open in $\overline{O(s_1, \dots, s_n)}$, since its complement is closed. Thus, both $\overline{O}(s_1, \dots, s_n)$ and $\overline{O(s_1, \dots, s_n)}$ contain $O(s_1, \dots, s_n)$ as a dense subset.*

proof: For notational simplicity, let $X = \mathcal{B}$, let $Y_i = O(s_i) \subset X \times X$, and let $f_i : X^{n+1} \rightarrow Y_i$ be the morphism $f_i(x_0, \dots, x_n) = (x_{i-1}, x_i)$, for $1 \leq i \leq n$. Let

$$Z = \{x \in X^{n+1} \mid f_i(x) \in Y_i, \forall 1 \leq i \leq n\}.$$

To prove the lemma, it suffices to prove

$$\overline{Z} = \{x \in X^{n+1} \mid f_i(x) \in \overline{Y_i}, \forall 1 \leq i \leq n\}. \quad (2)$$

Let $g : X^{n+1} \rightarrow Y_1 \times \dots \times Y_n$, so $g = (f_1, \dots, f_n)$. Since each f_i is proper (and hence closed), so is g . Then $Z = g^{-1}(Y_1 \times \dots \times Y_n)$. We have

$$g^{-1}(Y_1 \times \dots \times Y_n) \subset \overline{g^{-1}(Y_1 \times \dots \times Y_n)} \subset g^{-1}(\overline{Y_1 \times \dots \times Y_n}),$$

since g is continuous in the Zariski topology. Recall G is connected, so each $Y_i = O(s_i)$ is irreducible. Since g is also closed, it follows that $g^{-1}(Y_1 \times \dots \times Y_n)$ is open in its closure. This implies (2). $\square$

Next we determine the reduced fibers of π_n .

Proposition 2.3. *For $n \geq 1$, the projection morphism induces a surjective morphism*

$$\pi_n : \overline{O}(s_1, \dots, s_n) \rightarrow \overline{O}(s_1, \dots, s_{n-1}),$$

whose reduced fibers are isomorphic to $\mathbb{P}^1$.

Remark 2.4. (1) *The morphism π_n is proper, since it can be expressed as the composition of a closed immersion (inclusion) with a base-change morphism (projection $\mathcal{B}^{n+1} \rightarrow \mathcal{B}^n$).*

(2) *In the case when $w = s_1 \dots s_n$ is a minimal expression, see for example, [Spr1], §10.2.13.*

proof: We want to show the projection map

$$\overline{O}(s_1, \dots, s_n) \rightarrow \overline{O}(s_1, \dots, s_{n-1})$$

is surjective with fibers $\mathbb{P}^1$. Observe that there is an isomorphism of reduced fibers

$$\pi_n^{-1}(B_0, \dots, B_{n-1}) \rightarrow \pi_1^{-1}(B_{n-1}), \quad (3)$$

given by $(B'_0, \dots, B'_n) \mapsto (B'_{n-1}, B'_n)$. This boils the proof down to investigating

$$\pi_1^{-1}(B') = (B' \times \mathcal{B}) \cap \overline{O}(s) = \{(B', B'') \mid (B', B'') \in \overline{O(s)}\},$$

where B' is fixed and s is simple.

For α a root of B' , let U_α denote the unipotent subgroup of B' associated to α and T the maximal torus of B' . Note

$$B' = T \times \prod_{\alpha > 0} U_\alpha,$$

and

$${}^s B' = T \times \prod_{s(\alpha) > 0} U_\alpha,$$

where $>$ denotes the Bruhat ordering with respect to (B', T) . Let $P_s = \langle B', {}^s B' \rangle$ be the parabolic generated by B' and ${}^s B'$. For $(B', B'') \in \pi_1^{-1}(B')$, we have $B'' = pB'p^{-1}$, for some $p \in P_s$ (the conjugating element must belong to the parabolic generated by the two Borels by Corollary 11.17(iii) in [Bor]). Let $Y = P_s/B'$. By [St], §3.9, we have $Y \cong \mathbb{P}^1$. Since a Borel is its own normalizer, we have an isomorphism $Y \rightarrow \pi_1^{-1}(B')$. The result follows. $\square$

Corollary 2.5. $\overline{O}(s_1, \dots, s_n)$ is irreducible.

proof: We assume for the moment that π_n is flat (this will be proven in Lemma 3.1). Since G is connected, $\overline{O}(\emptyset) = \mathcal{B}$ is irreducible. Since the reduced fibers of π_1 are $\cong \mathbb{P}^1$ (see Proposition 2.3), it follows that $\overline{O}(s_1)$ is also irreducible, by Lemma 5.3 in Debarre [De]. The claim follows by induction. $\square$

Remark 2.6. By induction, it follows that $\dim(\overline{O}(s_1, \dots, s_n)) = n + \dim(\mathcal{B})$.

Next, we prove Lemma 1.1.

Lemma 2.7. Let $H_n = G \times P_1 \times P_2 \times \dots \times P_n$, where B is our fixed Borel, and $P_i = \langle B, {}^{s_i} B \rangle$, $1 \leq i \leq n$. Let H_n operate² on $\mathcal{B}^{n+1}$ by

$$(g, p_1, \dots, p_n) : (B_0, \dots, B_n) \mapsto ({}^g B_0, {}^{gp_1} B_1, \dots, {}^{gp_1 \dots p_n} B_n),$$

for $p_i \in P_i$, $1 \leq i \leq n$, and $g \in G$.

- (a) $\overline{O}(s_1, \dots, s_n)$ is the orbit of H_n operating on $x_{n+1} = (B, \dots, B) \in \mathcal{B}^{n+1}$: $\overline{O}(s_1, \dots, s_n) = H_n x_{n+1}$.
- (b) There is a projection $\rho_n : H_n \rightarrow H_{n-1}$ such that, for all $h \in H_n$, $\pi_n(h \cdot x_{n+1}) = \rho_n(h) \cdot \pi_n(x_{n+1}) = \rho_n(h) \cdot x_n$.
- (c) Consider the projection $pr_1 : \overline{O}(s_1, \dots, s_n) \rightarrow \mathcal{B}$, sending $(B_0, \dots, B_n) \mapsto B_0$. Then $pr_1^{-1}(B) = H'_n x_{n+1}$, where $H'_n = P_1 \times P_2 \times \dots \times P_n$ and the operation of H'_n is via the natural embedding $H'_n \hookrightarrow H_n$.

²This is not an action.

- (d) Let $h, h' \in H'_n$ be given by $h = (p_1, \dots, p_n)$ and $h' = (p'_1, \dots, p'_n)$, where $p_i, p'_i \in P_i$. We have $h \cdot x_{n+1} = h' \cdot x_{n+1}$ if and only if $p'_i p_i^{-1} \in {}^{p_{i-1} \dots p_1} B$, for all $1 \leq i \leq n$ (when $i = 1$, this should be interpreted as $p'_1 p_1^{-1} \in B$).
- (e) Define a relation $\sim$ on H'_n as follows: if $h, h' \in H'_n$ are given by $h = (p_1, \dots, p_n)$ and $h' = (p'_1, \dots, p'_n)$, where $p_i, p'_i \in P_i$, then define $h \sim h'$ if and only if $p'_i p_i^{-1} \in {}^{p_{i-1} \dots p_1} B$, for all $1 \leq i \leq n$ (when $i = 1$, this should be interpreted as $p'_1 p_1^{-1} \in B$). This is an equivalence relation.

$$pr_1^{-1}(B) \cong H'_n / \sim,$$

as sets.

Remark 2.8. (1) The proof below shows how to realize the Demazure varieties denoted Z_K in §3 of [D] as fibers in $\overline{O}(s_1, \dots, s_n)$.

(2) If, in addition, $s_1, s_1 s_2, \dots, s_1 \dots s_n$ is a Hamiltonian path in the Cayley graph of the Weyl group W generated by the simple reflections of B then this fact also explains how to embed Langlands' variety of stars S [L] into $\overline{O}(s_1, \dots, s_n)$.

(3) Note that Z_n (and, by Corollary 2.5, $\overline{O}(s_1, \dots, s_n)$) is irreducible, since G is connected.

proof: (a): Consider the H_n -orbit of x_{n+1} : $Z_n = H_n x_{n+1}$. We claim $Z_n = \overline{O}(s_1, \dots, s_n)$.

By construction, $Z_n \subset \overline{O}(s_1, \dots, s_n)$. The projection $\pi'_n : Z_n = H_n x_{n+1} \rightarrow Z_{n-1} = H_{n-1} x_n$ sends the point $({}^g B, {}^{gp_1} B, \dots, {}^{gp_1 \dots p_n} B) \in Z_n$ to $({}^g B, {}^{gp_1} B, \dots, {}^{gp_1 \dots p_{n-1}} B) \in Z_{n-1}$.

We identify the fiber of π'_n over $x_n \in Z_{n-1}$, $(\pi'_n)^{-1}(x_n)$. By definition,

$$(\pi'_n)^{-1}(x_n) = \{({}^g B, {}^{gp_1} B, \dots, {}^{gp_1 \dots p_n} B) \in Z_n \mid {}^g B = B, {}^{gp_1} B = B, \dots, {}^{gp_1 \dots p_{n-1}} B = B\}.$$

Since B is its own normalizer, these conditions become $g \in B, gp_1 \in B$ hence $p_1 \in B, \dots, gp_1 \dots p_{n-1} \in B$ hence $p_{n-1} \in B$. The only condition left unaffected then is that on p_n , so $p_n \in P_n/B$. We therefore see that this reduced fiber is $\cong P_n/B$, which is $\cong \mathbb{P}^1$ as in the proof of Proposition 2.3. Also, note that the projection π'_n is compatible with the inclusion $Z_n \subset \overline{O}(s_1, \dots, s_n)$. Thus the fiber of π'_n over x_n is contained in the corresponding fiber of π_n . As both of these fibers are isomorphic to $\mathbb{P}^1$, they must be equal. By the inclusion, each fiber of π'_n is contained in the corresponding fiber of π_n . We claim that

each fiber of p'_n is of the form $h \cdot (\pi'_n)^{-1}(x_n)$, for some $h \in H_n$. Thus all fibers of p'_n are isomorphic to $\mathbb{P}^1$. Thus the fiber of π'_n over any point is contained in the corresponding fiber of π_n . As both of these fibers are isomorphic to $\mathbb{P}^1$, they must be equal.

On the other hand, $Z_0 \cong G/B \cong \mathcal{B}$. Since the fibers of π_1 and π'_1 are equal, we have $Z_1 = \overline{O}(s_1)$. By induction, we have $Z_n = \overline{O}(s_1, \dots, s_n)$. This proves (a).

(b), (c): An easy consequence of (a).

(d): Suppose

$$(B, {}^{p_1}B, \dots, {}^{p_1 \dots p_n}B) = (B, {}^{p'_1}B, \dots, {}^{p'_1 \dots p'_n}B).$$

Since B is its own normalizer in G , ${}^{p_1}B = {}^{p'_1}B$ implies $p'_1 p_1^{-1} \in B$. Similarly, ${}^{p_1 p_2}B = {}^{p'_1 p'_2}B$ implies $p'_2 p_2^{-1} \in {}^{p_1}B$. The claim in the lemma follows by an induction argument.

(e): To check that $\sim$ is an equivalence relation, note that the condition

$$p_i(p'_i)^{-1} \in {}^{p_{i-1} \dots p_1}B, \quad \forall 1 \leq i \leq n,$$

is equivalent to

$$(p_1 \dots p_i)(p'_1 \dots p'_i)^{-1} \in B, \quad \forall 1 \leq i \leq n.$$

Each of this last set of conditions is easily seen to be both symmetric and transitive.

The last claim of (e) follows from (d) and the fact that $\sim$ is an equivalence relation. $\square$

Proposition 2.9. *Assume that k has characteristic 0. Each schematic fiber of π_n is non-singular.*

We conjecture this holds in characteristic $p > 0$.

proof: First, we prove the above proposition in the case $n = 1$. The property of being “geometrically reduced” holds on an open set (Theorem 12.2.4 in ch IV of [EGA]). We will show that, as a consequence of more general results proven in section §6, this open set contains the generic point, hence is dense (and thus is non-empty). As there is a transitive group action of G on the set of these fibers (Lemma 2.7 (b)), all the fibers are isomorphic (as schemes), hence they must all be geometrically reduced. We conclude from Proposition 2.3 that these fibers are all (reduced schemes) isomorphic to $\mathbb{P}^1$.

Now the proposition follows from the schematic analog of (3), which is a special case of (5). $\square$

3 The smoothness result

We shall use the following result.

Lemma 3.1. *For each $(w_1, \dots, w_n) \in W^n$, we have*

$$\overline{O}(w_1, \dots, w_n) = \overline{O}(w_1, \dots, w_{n-1}) \times_{\mathcal{B}} \overline{O}(w_n),$$

where the morphism $\overline{O}(w_1, \dots, w_{n-1}) \rightarrow \mathcal{B}$ is given by $(B_0, \dots, B_{n-1}) \mapsto B_{n-1}$ and the morphism $\overline{O}(w_n) \rightarrow \mathcal{B}$ is given by $\pi_1 : (B'_0, B'_1) \mapsto B'_0$. The projection morphism π_n is flat.

proof: Let $X' = \overline{O}(w_1, \dots, w_n)$, $X = \overline{O}(w_1, \dots, w_{n-1})$, $Y' = \overline{O}(w_n)$, and let $Y = \mathcal{B}$. The fact that $X' = X \times_Y Y'$ follows from the definition of $X' = \overline{O}(w_1, \dots, w_n)$ and $\times_Y$. We must show that $\pi_1 : Y' \rightarrow Y$ is flat.

A well-known theorem implies that π_1 is generically flat (see Theorem 6.9.1 in ch IV of [EGA] or for example, Corollary III.10.7 in [H]). Since G acts transitively on Y and π_1 is G -equivariant by Lemma 2.7 (b), π_1 must be flat over $O(w_n) \subset Y'$.

Note that G acts transitively on the fibers of π_1 . In particular, given any two $x, x' \in \overline{O}(w_n)$ with, say $x \in \pi_1^{-1}(z)$ and $x' \in \pi_1^{-1}(z')$ for $z, z' \in \mathcal{B}$, there is a $g \in G$ such that $gz = z'$. In other words, we can “translate” any fiber of any point in $\overline{O}(w_n)$ to the fiber of a point in $O(w_n)$. Since π_1 is flat over $O(w_n)$, it follows π_1 is flat everywhere on $\overline{O}(w_n)$.

The conclusion that π_n is flat follows from “base change” (Proposition III.9.2(b) in Hartshorne [H]). $\square$

Remark 3.2. *To help visualize the geometry, we construct an open $U \subset Y'$ containing $Y' - O(w_n)$. Let B' denote a Borel opposite to B , let N' denote the unipotent radical of B' , and let $P_w = \langle B, {}^w B \rangle$ ($w \in W$). Define*

$$U(w) = \{({}^n B, {}^n p B) \mid n \in N', p \in P_w\}. \quad (4)$$

The map $N' \times P_{w_n}/B \rightarrow U(w_n)$, defined by $(n, p) \mapsto ({}^n B, {}^n p B)$, is an isomorphism.

Next we prove Proposition 1.2: for $\text{char}(k) = 0$, the closed variety of Borels of relative position $\vec{s}$ is non-singular and π_n is smooth.

proof: We prove this by induction on n , using Lemma 6.1.

Let $n = 1$. The reduced fibers are all $\mathbb{P}^1$ by Proposition 2.3. Since G acts transitively on $O(s_1)$, it must be non-singular. Each point in $U(s_1)$ is smooth,

where U is as in (4), so $\overline{O}(s_1)$ is non-singular. Each schematic fiber of π_1 is non-singular, by Proposition 2.9. Since each schematic fiber is non-singular, and its reduced scheme is $\mathbb{P}^1$, they must all be (isomorphic to) $\mathbb{P}^1$.

Since the schematic fibers are non-singular and π_1 is flat, by Theorem 3' in §III.10 of Mumford [Mum], π_1 is smooth.

Now assume $n > 1$. By the induction hypothesis, $\overline{O}(s_1, \dots, s_{n-1})$ is non-singular. As above, it suffices to check that π_n is flat (by Mumford's result and Proposition 2.9). So by Lemma 3.1 again, π_n is smooth. By the induction hypothesis and Lemma 6.1, $\overline{O}(s_1, \dots, s_n)$ is non-singular.

The schematic fibers are all $\mathbb{P}^1$ by the same argument as in the case $n = 1$. $\square$

4 Remarks on $\mathbb{P}^1$ -bundles

In this section, we discuss Question 1.4.

First, we recall the definition of a $\mathbb{P}^1$ -bundle. A $\mathbb{P}^1$ -**bundle** E over X is (1) a morphism $\varphi : E \rightarrow X$, (2) an open covering $\{U_\alpha\}$ of X and a collection of isomorphisms $\varphi_\alpha : \varphi^{-1}(U_\alpha) \rightarrow U_\alpha \times \mathbb{P}^1$ such that (a) the following diagram commutes

$$\begin{array}{ccc} \varphi^{-1}(U_\alpha) & \xrightarrow{\varphi_\alpha} & U_\alpha \times \mathbb{P}^1 \\ \varphi \downarrow & & \downarrow pr_1 \\ U_\alpha & \xlongequal{\quad} & U_\alpha, \end{array}$$

and (b) for all α, β ,

$$\varphi_\beta \varphi_\alpha^{-1} : U_\alpha \cap U_\beta \times \mathbb{P}^1 \rightarrow U_\alpha \cap U_\beta \times \mathbb{P}^1$$

may be regarded as an element of $\Gamma(U_\alpha \cap U_\beta, PGL(2))$. This definition is essentially a special case of that in Hartshorne [H], page 170.

Regarding the question of whether or not

$$\overline{O}(s_1, \dots, s_n) \xrightarrow{\pi_n} \overline{O}(s_1, \dots, s_{n-1})$$

is a $\mathbb{P}^1$ -bundle (so $\varphi = \pi_n$ in part (1) of the definition above), let us consider the case where $n = 1$: $\overline{O}(s) \xrightarrow{\pi_1} \overline{O}(\emptyset) = \mathcal{B}$. Write $(B_0, B_1) \in \overline{O}(s)$ as $B_0 = gB$, $B_1 = gsB$, for some simple reflection s (with respect to the Borel B) and some $g \in G$. Under π_1 , the pair $(B_0, B_1) \in \overline{O}(s)$ is sent to $B_0 \in \overline{O}(\emptyset)$.

The map $N' \times P_s/B \rightarrow U(s)$, defined by $(n, p) \mapsto ({}^n B, {}^n p B)$, is an isomorphism, where $U(s)$ is as in (4) and $P_s = \langle B, {}^s B \rangle$. Let $\psi_n : {}^n P_s / {}^n B \rightarrow \mathbb{P}^1$ denote an isomorphism. Consider the map $\phi : N' \times \mathbb{P}^1 \rightarrow \overline{O}(s)$ defined by

$$\begin{array}{ccc} N' \times \mathbb{P}^1 & \rightarrow & N' \times {}^n P_s / {}^n B \rightarrow \overline{O}(s) \\ (n, x) & \mapsto & (n, \psi_n(x)) \mapsto ({}^n B, {}^{n\psi(x)} B) \end{array}$$

In this case, the next step to addressing the question is to verify that the following diagram commutes:

$$\begin{array}{ccc} N' \times \mathbb{P}^1 & \xrightarrow{\phi} & \pi_1^{-1}(N') \\ pr_1 \downarrow & & \downarrow \pi_1 \\ N' & \xlongequal{\quad} & N'. \end{array}$$

This follows from the definition. Therefore, we have some evidence³ that condition (2a) of the definition above holds, in the case $n = 1$.

Although our discussion above does not settle the question, from what we have shown it follows that $\overline{O}(s)$ is a uniruled variety, in the sense of Debarre [De].

5 Deligne-Lusztig varieties

We prove Corollary 1.5, under the assumption that Proposition 1.2 holds for $k = \overline{\mathbb{F}}_q$.

Let $\vec{s} = (s_1, \dots, s_n) \in S^n$ be simple reflections with respect to a fixed Borel B in a connected semisimple group G defined over a finite field $\mathbb{F}_q$. Let $\overline{X}(\vec{s})$ be the Deligne-Lusztig variety, as in §1. From [DL], page 151, we know that this variety is given by the fiber product

$$\overline{X}(\vec{s}) = \overline{O}(\vec{s}) \times_{B \times B} \Gamma,$$

where Γ is the graph of the Frobenius morphism F . By Lemma 6.1, Proposition 1.2, and the fact that the intersection is transversal (see [DL], Lemma 9.11), $\overline{X}(\vec{s})$ is non-singular. $\square$

³We have not even shown, among other things, that the above map ϕ represents a morphism.

6 Appendix: Technical background

This section is devoted to finishing the proof of Proposition 2.9.

First, we record a simple observation used repeatedly in the previous sections.

Lemma 6.1. *Let $f : X \rightarrow Y$ be a smooth morphism of varieties defined over k . If Y is non-singular then X is also non-singular.*

This is well-known.

6.1 Standard properties of fibers

We recall some standard facts due to Grothendieck [EGA].

From EGA I, 3.6.3: Assume that X and Y are affine schemes, that $F : X \rightarrow Y$ is a morphism, and that $U \subset X$ is open. The following diagram is commutative.

$$\begin{array}{ccc} X & \longleftarrow & U \\ f \downarrow & & \downarrow f|_U \\ Y & \xlongequal{\quad} & Y, \end{array}$$

where the top arrow is inclusion. For $y \in f(U)$, the morphism

$$f^{-1}(U) \cap U \rightarrow (f|_U)^{-1}(y)$$

induced from f , is an isomorphism.

From EGA I, 3.6.4: Let Y' be an irreducible component of Y and let $f : X \rightarrow Y$. Let

$$X_{(Y')} = X \times_Y Y' \xrightarrow{f_{(Y')}} Y'.$$

We have $f^{-1}(Y') = X' = X_{(Y')}$. Moreover,

$$\begin{array}{ccc} X_{(Y')} & \longrightarrow & X \\ f_{(Y')} \downarrow & & \downarrow f \\ Y' & \xrightarrow{j} & Y, \end{array}$$

where j is inclusion. Then

$$f^{-1}(y) \leftarrow f_{(Y')}^{-1}(y'), \tag{5}$$

is an isomorphism as schemes (“transitivity of fibers”).

6.2 Main lemma

Here is our main lemma.

Lemma 6.2. *Let D be an integral domain with field of fractions, $K = \text{Frac}(D)$. Let A be a D -algebra, flat over D and let $A_K = A \otimes_D K$. Then*

(i) *if A is an integral domain then so is A_K ,*

(ii) *if A is reduced then so is A_K .*

Remark 6.3. *Here is an example to show that the extension of (b) to “if A is reduced then A_K is geometrically reduced” is false. (This counterexample suggests that the “ $\text{char}(k) = 0$ ” hypothesis in Proposition 6.4 below cannot be removed, at least not without adding some other condition.) Let $k = \overline{\mathbb{F}_2}$, $D = k[x]$, $K = k(x)$, so $A_K = k(x)[y]/(y^2 - x)$. Consider the purely inseparable extension $K = k(\sqrt{x})$ of $K = k(x)$. Then $A_{K'} = k(\sqrt{x})[y]/(y^2 - x)$. Consider the non-zero element $a = y + \sqrt{x} \in A_{K'}$. This element satisfies $a^2 = 0$, so A_K is not geometrically reduced.*

proof: By flatness, the map

$$A = A \otimes_D D \rightarrow A_K$$

is injective, so we may view A as a subset of A_K .

For (i): Assume A_K is not a domain and assume, that $(\sum x \otimes y) \cdot (\sum x' \otimes y') = 0$, for some non-zero $\sum x \otimes y, \sum x' \otimes y' \in A_K$. Eliminating denominators, we may assume $y, y' \in D$. This gives divisors of zero in $A \otimes_D D = A \subset A_K$. This proves (i).

The proof of (ii) is similar, hence omitted.

□

Proposition 6.4. *Assume that k has characteristic 0. Let X and Y be reduced schemes defined over k and let $f : X \rightarrow Y$ be a proper, surjective, flat morphism of finite presentation. Then*

(i) *The generic fiber of f is geometrically reduced.*

(ii) *The general fiber of f is geometrically reduced (i.e., all fibers over a dense open set in Y are reduced).*

Remark 6.5. *We only need to assume f is flat for part (i).*

proof: We only prove the generic fiber of f is reduced here. The next subsection extends the proof to the geometrically reduced situation.

Part (ii) follows from (i) and [EGA] IV, §12.2.4.

It remains to prove (i). Using the standard properties of fibers (in §6.1) and the fact that Y is Noetherian, we reduce to the case where X and Y are affine and Y is integral. Appealing to Lemma 6.2 proves (i) with “geometrically reduced” replaced by “reduced”.

It remains to show that the generic fiber of f is geometrically reduced. By another result of Grothendieck, it suffices to show the fiber remains reduced after passing to a finite purely inseparable extension. But if k is perfect then every finite purely inseparable extension is trivial. $\square$

References

- [Bor] A. Borel, **Linear algebraic groups**, 2nd ed, Springer-Verlag, 1991
- [DL] P. Deligne and G. Lusztig, “Representations of reductive groups over finite fields,” *Ann. Math.* 103(1976)103-161
- [De] O. Debarre, **Higher-dimensional algebraic geometry**, Springer-Verlag, 2001
- [D] M. Demazure, “Désingularisation des variétés Schubert généralisées”, *Ann. Sci. École Norm. Sup.* 7(1974)53-88
- [EGA] J. Dieudonné and A. Grothendieck, “Eléments de géométrie algébrique,” *IHES* 4(1960), 24(1965), 28(1965)
- [Han1] S. H. Hansen, “Canonical bundles of Deligne-Lusztig varieties,” *Manu. Math.* 98(1999)363-375
- [Han2] —, “Error-correcting codes from higher dimensional varieties,” *Finite Fields and Appl.* 7(2001)530-552
- [H] R. Hartshorne, **Algebraic geometry**, Springer-Verlag, 1977.
- [L] R. Langlands, “Orbital integrals on forms of $SL(3)$, I”, *Amer J Math* 105(1983) 465-506

- [Mum] D. Mumford, **The red book of varieties and schemes**, Springer-Verlag LNM 1358, 1988
- [Spr1] T. A. Springer, **Linear algebraic groups**, 1st ed., Birkäuser, Boston, 1983
- [Spr2] ———, **Linear algebraic groups**, 2nd ed., Birkäuser, Boston, 1998
- [St] R. Steinberg, **Conjugacy classes in algebraic groups**, Springer LNM 366, 1974